


Electrodynamics

— Lecture B —

UNIVERSITY
OF TWENTE.

MESA+
INSTITUTE FOR NANOTECHNOLOGY



Manfred Hammer*, Hugo Hoekstra,
Lantian Chang, Mustafa Sefünc, Yean-Sheng Yong

Integrated Optical MicroSystems
MESA⁺ Institute for Nanotechnology
University of Twente, The Netherlands

University of Twente, Enschede, The Netherlands — Course 191210410(1), Block 1A, 2013/2014

* Department of Electrical Engineering, University of Twente
Phone: +31/53/489-3448

P.O. Box 217, 7500 AE Enschede, The Netherlands
Fax: +31/53/489-3996

E-mail: m.hammer@utwente.nl

Course overview

Electrodynamics

- A Introduction, Maxwell equations, brush up on vector calculus, Dirac delta, potentials, Taylor expansion, Fourier transform,
- B Maxwell equations, brush up on electro- and magnetostatics, multipole expansion
- C Maxwell equations, microscopic and macroscopic, time- and frequency domain, differential and integral form, interface conditions, continuity equation, energy and momentum of electromagnetic fields
- D Wave equation, plane waves, plane harmonic electromagnetic waves, refractive index, polarization, energy transport, spherical waves
- E Reflection and transmission at interfaces, lossy materials, wave packets, dispersion, phase and group velocity
- F Maxwell equations, vectorial and scalar Helmholtz equation, 2D configurations (intermezzo), mode problems, metallic and dielectric waveguides
- G Scalar and vector potentials, gauge conditions, retarded potentials, electric and magnetic dipole radiation
- H Special relativity; transmission lines

Maxwell equations

SI, in matter, time domain, differential form:

$\nabla \cdot \mathbf{D} = \rho_f,$	$\mathbf{E}(\mathbf{r}, t)$: electric field,
$\nabla \times \mathbf{E} = -\dot{\mathbf{B}},$	$\mathbf{D}(\mathbf{r}, t)$: (di-)electric displacement,
$\nabla \cdot \mathbf{B} = 0,$	$\mathbf{B}(\mathbf{r}, t)$: magnetic induction (field, flux density),
$\nabla \times \mathbf{H} = \mathbf{J}_f + \dot{\mathbf{D}},$	$\mathbf{H}(\mathbf{r}, t)$: magnetic field (...),
$\mathbf{D} = \epsilon_0 \mathbf{E} + \mathbf{P},$	$\rho_f(\mathbf{r}, t)$: density of free charges,
$\mathbf{B} = \mu_0(\mathbf{H} + \mathbf{M}).$	$\mathbf{J}_f(\mathbf{r}, t)$: density of free currents,
	$\mathbf{P}(\mathbf{r}, t)$: polarization,
	$\mathbf{M}(\mathbf{r}, t)$: magnetization,
	ϵ_0 : free space permittivity,
(+ constitutive relations)	μ_0 : free space permeability.

$\mathbf{F} = q(\mathbf{E} + \mathbf{v} \times \mathbf{B})$ (Lorentz-) force \mathbf{F} on a charge q with velocity \mathbf{v} .

Polarization

\mathbf{P} : density of electric dipole moment (bound charges).

$$\mathbf{D} = \epsilon_0 \mathbf{E} + \mathbf{P}, \quad [\mathbf{D}] = [\mathbf{P}] = \frac{\text{As m}}{\text{m}^3}, \quad [\mathbf{E}] = \frac{\text{V}}{\text{m}},$$

vacuum permittivity $\epsilon_0 = 8.854187817 \dots \cdot 10^{-12} \left[\frac{\text{F}}{\text{m}} = \frac{\text{As}}{\text{Vm}} \right]$.

- Local dipoles induced by \mathbf{E}  $\mathbf{P}(\mathbf{E})$.

- Simplest case: linear isotropic dielectrics,

$$\begin{aligned} \mathbf{P} &= \epsilon_0 \chi_e \mathbf{E}, & \chi_e: \text{dielectric susceptibility, } [\chi_e] &= 1. \\ \mathbf{D} &= \epsilon_0 (1 + \chi_e) \mathbf{E} = \epsilon_0 \epsilon \mathbf{E}, & \epsilon: \text{relative permittivity, } [\epsilon] &= 1. \end{aligned}$$

- Complications: $\epsilon(\mathbf{r})$, $\epsilon(\omega)$, ϵ_{jk} , $\text{Im}\epsilon$, $\epsilon(T)$, $\epsilon(\mathbf{F})$, $\chi_{jkl}^{(2)} E_k E_l$, $\chi_{jklm}^{(3)} E_k E_l E_m$, \dots

Magnetization

M : density of magnetic dipole moments (bound currents).

$$\mathbf{H} = \frac{1}{\mu_0} \mathbf{B} - \mathbf{M}, \quad [\mathbf{H}] = [\mathbf{M}] = \frac{\text{A m}^2}{\text{m}^3}, \quad [\mathbf{B}] = \text{T} = \frac{\text{Vs}}{\text{m}^2},$$

$$\text{vacuum permeability } \mu_0 = 4\pi \cdot 10^{-7} \left[\frac{\text{N}}{\text{A}^2} = \frac{\text{Vs}}{\text{Am}} \right].$$

- Local dipoles induced by \mathbf{H}  $\mathbf{M}(\mathbf{H})$.

- Simplest case: linear isotropic magnetic media,

$$\begin{aligned} \mathbf{M} &= \chi_m \mathbf{H}, & \chi_m: \text{magnetic susceptibility, } [\chi_m] &= 1. \\ \mathbf{B} &= \mu_0(1 + \chi_m) \mathbf{H} = \mu_0 \mu \mathbf{H}, & \mu: \text{relative permeability, } [\mu] &= 1. \end{aligned}$$

- Complications: manifold.

- $$\begin{aligned}\nabla \cdot \boldsymbol{D} &= \rho_{\text{f}}, & \nabla \times \boldsymbol{E} &= -\dot{\boldsymbol{B}}, & \boldsymbol{D} &= \epsilon_0 \epsilon \boldsymbol{E}, \\ \nabla \cdot \boldsymbol{B} &= 0, & \nabla \times \boldsymbol{H} &= \boldsymbol{J}_{\text{f}} + \dot{\boldsymbol{D}}, & \boldsymbol{B} &= \mu_0 \mu \boldsymbol{H}.\end{aligned}$$

-  Magnetostatics.

- $$\nabla \cdot \mathbf{B} = 0, \quad \nabla \times \mathbf{H} = \mathbf{J}_f, \quad \mathbf{B} = \mu_0 \mu \mathbf{H} \quad \rightsquigarrow \text{Induction, AC currents, ...}$$

Electrostatics

$$\nabla \cdot \mathbf{D} = \rho_f, \quad \nabla \times \mathbf{E} = 0, \quad \mathbf{D} = \epsilon_0 \epsilon \mathbf{E}$$

↪ $\nabla \cdot (\epsilon \mathbf{E}) = \frac{\rho_f}{\epsilon_0}, \quad \nabla \times \mathbf{E} = 0$ $\epsilon(\mathbf{r}) !!!$

- Homogeneous media (piecewise ...), $\partial_r \epsilon = \epsilon \partial_r$:

$$\nabla \cdot \mathbf{E} = \frac{\rho_f}{\epsilon_0 \epsilon}, \quad \nabla \times \mathbf{E} = 0.$$

- Physical Gauss' law:

$Q_{f,\mathcal{V}}$: free charges in \mathcal{V}

$$\int_{\partial \mathcal{V}} \mathbf{E} \cdot d\mathbf{a} = \frac{Q_{f,\mathcal{V}}}{\epsilon \epsilon_0}.$$

- Scalar electrostatic potential $\phi(\mathbf{r})$:

(lecture A, potentials ...)

$$\mathbf{E} = -\nabla \phi, \quad \phi(\mathbf{r}) - \phi(\mathbf{r}_0) = - \int_{\mathbf{r}_0}^{\mathbf{r}} \mathbf{E}(\mathbf{r}') \cdot d\mathbf{r}',$$

- Poisson equation: $\Delta \phi = -\frac{\rho_f}{\epsilon_0 \epsilon}.$

Electrostatics, solutions

- Poisson equation: $\Delta\phi = -\frac{\rho_f}{\epsilon_0\epsilon}$.

- Solutions ($\mathcal{V} \rightarrow \infty$):

$$\phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0\epsilon} \int \frac{\rho_f(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathcal{V}', \quad \mathbf{E}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0\epsilon} \int \rho_f(\mathbf{r}') \frac{(\mathbf{r} - \mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|^3} d\mathcal{V}'.$$

- Point charge q at \mathbf{r}_0 , $\rho_f(\mathbf{r}) = q \delta(\mathbf{r} - \mathbf{r}_0)$:

$$\phi(\mathbf{r}) = \frac{q}{4\pi\epsilon_0\epsilon} \frac{1}{|\mathbf{r} - \mathbf{r}_0|}, \quad \mathbf{E}(\mathbf{r}) = \frac{q}{4\pi\epsilon_0\epsilon} \frac{(\mathbf{r} - \mathbf{r}_0)}{|\mathbf{r} - \mathbf{r}_0|^3}.$$

- Coulomb's law, force on another charge \tilde{q} at \mathbf{r} :

$$\mathbf{F}_{q \rightarrow \tilde{q}} = \frac{\tilde{q} q}{4\pi\epsilon_0\epsilon} \frac{(\mathbf{r} - \mathbf{r}_0)}{|\mathbf{r} - \mathbf{r}_0|^3}.$$

Electrostatics, charge densities

Variants of charge densities:

- $\rho(\mathbf{r})$, volume, $Q = \int_{\mathcal{V}} \rho \, d\mathcal{V}$, $[\rho] = \text{As}/m^3$,
- $\sigma(\mathbf{r})$, surface, $Q = \int_{\mathcal{S}} \sigma \, da$, $[\sigma] = \text{As}/m^2$,
- $\lambda(\mathbf{r})$, line, $Q = \int_{\mathcal{P}} \lambda \, ds$, $[\lambda] = \text{As}/m$,
- Q , point at \mathbf{r}_0 , $\rho(\mathbf{r}) = Q \delta(\mathbf{r} - \mathbf{r}_0)$, $[Q] = \text{As} = \text{C}$.

Electrostatics, conductors

Conductors, **electrostatic** configurations, free mobility of charges:

- External field induces charges σ_f **on the surface** of the conductor.
- These cancel the external field in the interior
 $\rightsquigarrow \mathbf{E} = 0$ **inside** the conductor.
- $\nabla \cdot \mathbf{E} = \frac{\rho_f}{\epsilon_0 \epsilon} \rightsquigarrow \rho_f = 0$ **inside** the conductor.
- $\phi(\mathbf{r}) = \phi(\mathbf{r}_0) - \int_{\mathbf{r}_0}^{\mathbf{r}} \mathbf{E} \cdot d\mathbf{s} \rightsquigarrow \phi$ constant over the conductor.
- $\mathbf{E}_{\text{outside}} \perp$ conductor surface.
- (Faraday cage).

Magnetostatics

$$\nabla \cdot \mathbf{B} = 0, \quad \nabla \times \mathbf{H} = \mathbf{J}_f, \quad \mathbf{B} = \mu_0 \mu \mathbf{H}$$


$$\nabla \cdot \mathbf{B} = 0, \quad \frac{1}{\mu_0} \nabla \times \frac{1}{\mu} \mathbf{B} = \mathbf{J}_f \qquad \mu(\mathbf{r}) !!!$$

- Homogeneous media (piecewise ...), $\partial_r \mu = \mu \partial_r$:

$$\nabla \cdot \mathbf{B} = 0, \quad \nabla \times \mathbf{B} = \mu_0 \mu \mathbf{J}_f.$$

- Ampère's circuital law:

$I_{f,\mathcal{S}}$: free current through \mathcal{S}

$$\oint_{\partial \mathcal{S}} \mathbf{B} \cdot d\mathbf{s} = \mu \mu_0 I_{f,\mathcal{S}}.$$

- Vector potential $\mathbf{A}(\mathbf{r})$:

(lecture A, potentials ...)

$$\mathbf{B} = \nabla \times \mathbf{A}.$$

- Poisson equation, vectorial: $\Delta \mathbf{A} = -\mu_0 \mu \mathbf{J}_f.$

Magnetostatics, solutions

- Poisson equation, vectorial: $\Delta \mathbf{A} = -\mu_0 \mu \mathbf{J}_f$.

- Solutions ($\mathcal{V} \rightarrow \infty$):

$$\mathbf{A}(\mathbf{r}) = \frac{\mu_0 \mu}{4\pi} \int \frac{\mathbf{J}_f(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathcal{V}', \quad \mathbf{B}(\mathbf{r}) = \frac{\mu_0 \mu}{4\pi} \int \mathbf{J}_f(\mathbf{r}') \times \frac{(\mathbf{r} - \mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|^3} d\mathcal{V}'.$$

- Current I along \mathcal{C} :

$$\mathbf{A}(\mathbf{r}) = \frac{\mu_0 \mu}{4\pi} I \oint_{\mathcal{C}} \frac{1}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{s}', \quad \mathbf{B}(\mathbf{r}) = \frac{\mu_0 \mu}{4\pi} I \oint_{\mathcal{C}} d\mathbf{s}' \times \frac{(\mathbf{r} - \mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|^3}.$$

- Ampère's force law, force on another current \tilde{I} along $\tilde{\mathcal{C}}$:

$$\mathbf{F}_{\mathcal{C} \rightarrow \tilde{\mathcal{C}}} = \frac{\mu_0 \mu}{4\pi} \tilde{I} I \oint_{\tilde{\mathcal{C}}} \oint_{\mathcal{C}} \frac{d\mathbf{s} \times (d\mathbf{s}' \times (\mathbf{r} - \mathbf{r}'))}{|\mathbf{r} - \mathbf{r}'|^3}.$$

Magnetostatics, current densities

Variants of current densities:

- $\mathbf{J}(\mathbf{r})$, volume, $I = \int_{\mathcal{S}} \mathbf{J} \cdot d\mathbf{a}$, $[\mathbf{J}] = \text{A}/\text{m}^2$,
- $\mathbf{K}(\mathbf{r})$, surface, $I = \int_{\mathcal{P}} \mathbf{K} \cdot \mathbf{t} ds$, $|\mathbf{t}| = 1$, $\mathbf{t} \parallel \text{surface}$, $\mathbf{t} \perp d\mathbf{s}$, $[\mathbf{K}] = \text{A}/\text{m}$,
- $\mathbf{I}(\mathbf{r})$, line, $I = |\mathbf{I}|$, $[\mathbf{I}] = \text{A}$,
- Current density \longleftrightarrow moving charge density:
 $\mathbf{J} = \rho \mathbf{v}$, $\mathbf{K} = \sigma \mathbf{v}$, $\mathbf{I} = \lambda \mathbf{v}$.

Multipole expansion

Electrostatics:

localized charge density $\rho_f(\mathbf{r})$, $\Delta\phi = -\frac{\rho_f}{\epsilon_0\epsilon}$,

↪
$$\phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0\epsilon} \int \frac{\rho_f(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathcal{V}'.$$

Magnetostatics:

localized current density $\mathbf{J}_f(\mathbf{r})$, $\Delta\mathbf{A} = -\mu_0\mu\mathbf{J}_f$,

↪
$$\mathbf{A}(\mathbf{r}) = \frac{\mu_0\mu}{4\pi} \int \frac{\mathbf{J}_f(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathcal{V}'.$$

Consider $\phi(\mathbf{r})$, $\mathbf{A}(\mathbf{r})$, for $|\mathbf{r}| \gg |\mathbf{r}'|_{\rho(r') \neq 0, J(r') \neq 0}$: ...

Multipole expansion, 1/r

Constant \mathbf{r}_s ; second order expansion of $\frac{1}{|\mathbf{r} - \mathbf{r}_s|}$ around $\mathbf{r} = \mathbf{0}$:

$$\frac{1}{|\Delta\mathbf{r} - \mathbf{r}_s|} \approx \frac{1}{|\mathbf{r}_s|} + \frac{\mathbf{r}_s \cdot \Delta\mathbf{r}}{|\mathbf{r}_s|^3} + \frac{1}{2} \left\{ \frac{3(\Delta\mathbf{r} \cdot \mathbf{r}_s)^2 - \mathbf{r}_s^2 \Delta\mathbf{r}^2}{|\mathbf{r}_s|^5} \right\} \pm \dots$$

(exercise)

Expansion required for $|\mathbf{r}| \gg |\mathbf{r}'|$; substitute $\Delta\mathbf{r} \rightarrow \mathbf{r}'$, $r' = |\mathbf{r}'|$,
 $\mathbf{r}_s \rightarrow \mathbf{r}$, $r = |\mathbf{r}|$:

$$\frac{1}{|\mathbf{r} - \mathbf{r}'|} \approx \frac{1}{r} + \frac{\mathbf{r} \cdot \mathbf{r}'}{r^3} + \frac{1}{2} \left\{ \frac{3(\mathbf{r}' \cdot \mathbf{r})^2 - r^2 (\mathbf{r}')^2}{r^5} \right\} \pm \dots$$

Multipole expansion, ϕ

$$\phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0\epsilon} \int \frac{\rho_f(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathcal{V}'$$

&

$$\frac{1}{|\mathbf{r} - \mathbf{r}'|} \approx \frac{1}{r} + \frac{\mathbf{r} \cdot \mathbf{r}'}{r^3} + \frac{1}{2} \left(\frac{3(\mathbf{r}' \cdot \mathbf{r})^2 - r^2(\mathbf{r}')^2}{r^5} \right) \pm \dots$$



$$\phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0\epsilon} \left\{ \begin{aligned} &\frac{1}{r} \int \rho_f(\mathbf{r}') d^3r' \\ &+ \frac{1}{r^3} \mathbf{r} \cdot \int \rho_f(\mathbf{r}') \mathbf{r}' d^3r' \\ &+ \frac{1}{2r^5} \int \rho_f(\mathbf{r}') (3(\mathbf{r}' \cdot \mathbf{r})^2 - r^2(\mathbf{r}')^2) d^3r' \pm \dots \end{aligned} \right\}$$

Multipole moments, ϕ

$$\phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0\epsilon} \left\{ \frac{Q}{r} + \frac{\mathbf{p} \cdot \mathbf{r}}{r^3} + \frac{1}{2} \sum_{j,k=1}^3 Q_{j,k} \frac{r_j r_k}{r^5} \pm \dots \right\},$$

$$Q = \int \rho_f(\mathbf{r}') d^3r', \quad \text{total charge, monopole moment,}$$

$$\mathbf{p} = \int \rho_f(\mathbf{r}') \mathbf{r}' d^3r', \quad \text{electric dipole moment,}$$

$$Q_{jk} = \int \rho_f(\mathbf{r}') (3r'_j r'_k - (\mathbf{r}')^2 \delta_{jk}) d^3r', \quad \text{electric quadrupole moment.}$$

- $|\mathbf{r}| \gg |\mathbf{r}'|_{\rho(\mathbf{r}') \neq 0}$:
- $Q \neq 0$: monopole dominant; $\phi(\mathbf{r}) \approx \frac{1}{4\pi\epsilon_0\epsilon} \frac{q}{r}$,
 - $Q = 0, \mathbf{p} \neq 0$: dipole dominant; $\phi(\mathbf{r}) \approx \frac{1}{4\pi\epsilon_0\epsilon} \frac{\mathbf{p} \cdot \mathbf{r}}{r^3}$,
 - $Q = 0, \mathbf{p} = 0$: quadru- (...) -pole dominant.

$\mathbf{E} = -\nabla\phi$  corresponding electric fields.

Multipole expansion, A

$$A(\mathbf{r}) = \frac{\mu_0 \mu}{4\pi} \int \frac{\mathbf{J}_f(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} dV'$$

&

$$\frac{1}{|\mathbf{r} - \mathbf{r}'|} \approx \frac{1}{r} + \frac{\mathbf{r} \cdot \mathbf{r}'}{r^3} \pm \dots$$



$$A(\mathbf{r}) = \frac{\mu_0 \mu}{4\pi} \left\{ \begin{aligned} &\frac{1}{r} \int \mathbf{J}_f(\mathbf{r}') d^3 r' \\ &+ \frac{1}{r^3} \int \mathbf{J}_f(\mathbf{r}') (\mathbf{r} \cdot \mathbf{r}') d^3 r' \\ &\pm \dots \end{aligned} \right\}$$

Multipole moments, \mathbf{A}

$$\mathbf{A}(\mathbf{r}) = \frac{\mu_0 \mu}{4\pi} \left\{ 0 + \frac{\mathbf{m} \times \mathbf{r}}{r^3} \pm \dots \right\},$$

no magnetic **monopoles**,

$$\mathbf{m} = \frac{1}{2} \int \mathbf{r}' \times \mathbf{J}_f(\mathbf{r}') d^3 r', \quad \text{magnetic **dipole** moment.}$$

- $|\mathbf{r}| \gg |\mathbf{r}'|_{\rho(\mathbf{r}') \neq 0}$:
- $\mathbf{m} \neq 0$: dipole term dominant; $\mathbf{A}(\mathbf{r}) \approx \frac{\mu_0 \mu}{4\pi} \frac{\mathbf{m} \times \mathbf{r}}{r^3}$,
 - $\mathbf{m} = 0$: higher order moments become relevant.

$\mathbf{B} = \nabla \times \mathbf{A}$  corresponding magnetic fields.

Upcoming

Next lecture:

- Maxwell equations: free space and in media, time- and frequency domain, differential and integral form, interface conditions, continuity equation
- Energy and momentum of electromagnetic fields, Poynting theorem

